


PROSAN+ DETERGENT CHLORINE TABLETS

BLOOD SPILLS



Wear appropriate PPE: Gloves & Apron. When using solution, wear safety glasses with side shields in case of splashing.



To contain Blood Spill, cover with paper towels. Prepare the Prosan Plus solution as per the chart below using cold potable (tap) water. Carefully pour over towels. Leave for 3 minutes. Dispose of all the Spill material as Clinical Waste.



Clean the spill area with Neutral Detergent (Detergent Wipes).



Keep prepared solution for no longer than 24 hours. Dispose of residual solution into an appropriate drain, sluice or sink with running water.



Remove gloves and apron and dispose of as Clinical Waste then wash hands thoroughly.

DILUTION INSTRUCTIONS

FOR DISINFECTION OF (DRY/WET) BLOOD SPILLAGES – 10,000 PPM AVAILABLE CHLORINE			CONTACT TIME
Disinfection of Dry and Wet Blood Spillages	Prosan Plus Tablets	Mix 1 tablet per 100ml water Or 5 tablets per 500ml water Or 10 tablets per 1 Litre water	3 Minutes

HEALTH & SAFETY

DO NOT use directly on Vomit or Urine Spills

DO NOT breathe in fumes, gas, vapours, powder or mist

AVOID contact with Eyes and Skin

AVOID prolonged contact with metals.
TAKE CARE: Solution will bleach fabrics

ALWAYS dispose of used materials as Clinical Waste

ALWAYS make a fresh solution every 24 hours

ALWAYS work in a well-ventilated area

ALWAYS replace tablet pot lid and store in a secure, dry place

USE THIS BIOCIDES SAFELY: REFER TO LOCAL HOSPITAL POLICY. REFER TO THE PRODUCT SAFETY DATA SHEET. READ THE PRODUCT LABEL.